


# Many Billions of Rocky Planets in the Habitable Zones around Red Dwarfs in the Milky Way

By European Southern Observatory 17 hours 9 minutes ago



PR Image eso1214a Artist's impression of sunset on the super-Earth world Gliese 667 Cc

A new result from ESO's HARPS planet finder shows that rocky planets not much bigger than Earth are very common in the habitable zones around faint red stars. The international team estimates that there are tens of billions of such planets in the Milky Way galaxy alone, and probably about one hundred in the Sun's immediate neighbourhood. This is the first direct measurement of the frequency of super-

Earths around red dwarfs, which account for 80% of the stars in the Milky Way.

This first direct estimate of the number of light planets around red dwarf stars has just been announced by an international team using observations with the HARPS spectrograph on the 3.6-metre telescope at ESO's La Silla Observatory in Chile [\[1\]](#). A recent announcement ([eso1204](#)), showing that planets are ubiquitous in our galaxy used a different method that was not sensitive to this important class of exoplanets.

The HARPS team has been searching for exoplanets orbiting the most common kind of star in the Milky Way — red dwarf stars (also known as M dwarfs [\[2\]](#)). These stars are faint and cool compared to the Sun, but very common and long-lived, and therefore account for 80% of all the stars in the Milky Way.

“Our new observations with HARPS mean that about 40% of all red dwarf stars have a super-Earth orbiting in the habitable zone where liquid water can exist on the surface of the planet,” says Xavier Bonfils (IPAG, Observatoire des Sciences de l'Univers de Grenoble, France), the leader of the team. “Because red dwarfs are so common — there are about 160 billion of them in the Milky Way — this leads us to the astonishing result that there are tens of billions of these planets in our galaxy alone.”

The HARPS team surveyed a carefully chosen sample of 102 red dwarf stars in the southern skies over a six-year period. A total of nine super-Earths (planets with masses between one and ten times that of Earth) were found, including two inside the habitable zones of Gliese 581 ([eso0915](#)) and Gliese 667 C respectively. The astronomers could estimate how heavy the planets were and how far from their stars they orbited.

By combining all the data, including observations of stars that did not have planets, and looking at the

fraction of existing planets that could be discovered, the team has been able to work out how common different sorts of planets are around red dwarfs. They find that the frequency of occurrence of super-Earths [3] in the habitable zone is 41% with a range from 28% to 95%.

On the other hand, more massive planets, similar to Jupiter and Saturn in our Solar System, are found to be rare around red dwarfs. Less than 12% of red dwarfs are expected to have giant planets (with masses between 100 and 1000 times that of the Earth).

As there are many red dwarf stars close to the Sun the new estimate means that there are probably about one hundred super-Earth planets in the habitable zones around stars in the neighbourhood of the Sun at distances less than about 30 light-years [4].

"The habitable zone around a red dwarf, where the temperature is suitable for liquid water to exist on the surface, is much closer to the star than the Earth is to the Sun," says Stéphane Udry (Geneva Observatory and member of the team). "But red dwarfs are known to be subject to stellar eruptions or flares, which may bathe the planet in X-rays or ultraviolet radiation, and which may make life there less likely."

One of the planets discovered in the HARPS survey of red dwarfs is Gliese 667 Cc [5]. This is the second planet in this triple star system (see [eso0939](#) for the first) and seems to be situated close to the centre of the habitable zone. Although this planet is more than four times heavier than the Earth it is the closest twin to Earth found so far and almost certainly has the right conditions for the existence of liquid water on its surface. This is the second super-Earth planet inside the habitable zone of a red dwarf discovered during this HARPS survey, after Gliese 581d was announced in 2007 and confirmed in 2009.

"Now that we know that there are many super-Earths around nearby red dwarfs we need to identify more of them using both HARPS and future instruments. Some of these planets are expected to pass in front of their parent star as they orbit — this will open up the exciting possibility of studying the planet's atmosphere and searching for signs of life," concludes Xavier Delfosse, another member of the team ([eso1210](#)).

## Notes

[1] HARPS measures the radial velocity of a star with extraordinary precision. A planet in orbit around a star causes the star to regularly move towards and away from a distant observer on Earth. Due to the Doppler effect, this radial velocity change induces a shift of the star's spectrum towards longer wavelengths as it moves away (called a redshift) and a blueshift (towards shorter wavelengths) as it approaches. This tiny shift of the star's spectrum can be measured with a high-precision spectrograph such as HARPS and used to infer the presence of a planet.

[2] These stars are called M dwarfs because they have the spectral class M. This is the coolest of the seven classes in the simplest scheme for classifying stars accordingly to decreasing temperature and the appearance of their spectra.

[3] Planets with a mass between one and ten times that of the Earth are called super-Earths. There are no such planets in our Solar System, but they appear to be very common around other stars. Discoveries of such planets in the habitable zones around their stars are very exciting because — if the planet were rocky and had water, like Earth — they could potentially be an abode of life.

[4] The astronomers used ten parsecs as their definition of "close". This corresponds to about 32.6 light-years.

[5] The name means that the planet is the second discovered (c) orbiting the third component (C) of the triple star system called Gliese 667. The bright stellar companions Gliese 667 A and B would be prominent in the skies of Gliese 667 Cc. The discovery of Gliese 667 Cc was independently announced by Guillem Anglada-Escude and colleagues in February 2012, roughly two months after the electronic preprint of the Bonfils et al. paper went online. This confirmation of the planets Gliese 667 Cb and Cc by Anglada-Escude and collaborators was largely based on HARPS observations and data processing of the European team that were made publicly available through the ESO archive.

## More information

This research was presented in a paper “The HARPS search for southern extra-solar planets XXXI. The M-dwarf sample”, by Bonfils et al. to appear in the journal *Astronomy & Astrophysics*.

The team is composed of X. Bonfils (UJF-Grenoble 1 / CNRS-INSU, Institut de Planétologie et d’Astrophysique de Grenoble, France [IPAG]; Geneva Observatory, Switzerland), X. Delfosse (IPAG), S. Udry (Geneva Observatory), T. Forveille (IPAG), M. Mayor (Geneva Observatory), C. Perrier (IPAG), F. Bouchy (Institut d’Astrophysique de Paris, CNRS, France; Observatoire de Haute-Provence, France), M. Gillon (Université de Liège, Belgium; Geneva Observatory), C. Lovis (Geneva Observatory), F. Pepe (Geneva Observatory), D. Queloz (Geneva Observatory), N. C. Santos (Centro de Astrofísica da Universidade do Porto, Portugal), D. Ségransan (Geneva Observatory), J.-L. Bertaux (Service d’Aéronomie du CNRS, Verrières-le-Buisson, France), and Vasco Neves (Centro de Astrofísica da Universidade do Porto, Portugal and UJF-Grenoble 1 / CNRS-INSU, Institut de Planétologie et d’Astrophysique de Grenoble, France [IPAG]).

The year 2012 marks the 50th anniversary of the founding of the European Southern Observatory (ESO). ESO is the foremost intergovernmental astronomy organisation in Europe and the world’s most productive astronomical observatory. It is supported by 15 countries: Austria, Belgium, Brazil, the Czech Republic, Denmark, France, Finland, Germany, Italy, the Netherlands, Portugal, Spain, Sweden, Switzerland and the United Kingdom. ESO carries out an ambitious programme focused on the design, construction and operation of powerful ground-based observing facilities enabling astronomers to make important scientific discoveries. ESO also plays a leading role in promoting and organising cooperation in astronomical research. ESO operates three unique world-class observing sites in Chile: La Silla, Paranal and Chajnantor. At Paranal, ESO operates the Very Large Telescope, the world’s most advanced visible-light astronomical observatory and two survey telescopes. VISTA works in the infrared and is the world’s largest survey telescope and the VLT Survey Telescope is the largest telescope designed to exclusively survey the skies in visible light. ESO is the European partner of a revolutionary astronomical telescope ALMA, the largest astronomical project in existence. ESO is currently planning a 40-metre-class European Extremely Large optical/near-infrared Telescope, the E-ELT, which will become “the world’s biggest eye on the sky”.

Source: [European Southern Observatory](#)

Like

**Add New Comment**

[Login](#)